

To Remove

Carefully lift away the dressing and discard.

Will depend upon the amount of exudate. Adaptic may be left

Frequency of Dressing Change

in place for up to 7 days. **Expected Outcome**











To avoid trauma to the wound bed.





Skin and Wound Product Information Sheet

Skin and Wound Product Information Sheet		
Adaptic		
Classification	Layer: Petrolatum Impregnated	
Key Points	Petrolatum impregnated cellulose acetate dressing	
Indications	For prevention of wound bed trauma by decreasing adherence of the secondary dressing	
Precautions	Avoid using this product with silicone-type products as they have the same functionality	
Contraindications	 Do not use for clients with sensitivity or allergy to petrolatum Do not use during Hyperbaric Oxygen therapy procedure 	
Formats & Sizes	● Sheet ■ 7.6 x 7.6 cm ■ 7.6 x 20.3 cm ■ 7.6 x 40.6 cm ■ 12.7 x 22.9 cm	ARACHETE TO THE TOTAL THE
Application Directions		Rationale
Cleanse/irrigate wound with sterile normal saline or agency approved wound cleanser; dry peri-wound skin.		Reduces wound debris and allows for adhesion of dressing or tape.
If required, apply skin barrier to peri-wound skin.		To protect the peri-wound skin from maceration and to improve the adhesion of the dressing or tape.
To Apply		
Adaptic can be cut to size. Choose a size that covers the wound		To ensure adequate adhesion to peri-wound skin.
plus at least 2 cm of peri-wound skin.		
Apply Adaptic in a single layer over the wound and smooth onto the surrounding skin to get a good seal.		Single layer allows passage of exudate through to cover dressing.
Apply appropriate cover dressing to maintain a moisture-balanced wound environment.		The choice of the cover dressing will depend upon the amount of exudate expected.

Contact layer does not adhere to wound.

For further information, please contact your Wound Clinician.