







Product Information Sheet

Adapt Adhesive Remover		
Classification	Adhesive Remover	
British Columbia Practice	 The British Columbia Perinatal Services and the British Columbia Provincial Nursing Skin & Wound Committee have determined Adapt Adhesive Remover for neonates in: NICUs with a corrected gestational age greater than 31 weeks + 6 days. Maternity/pediatric units who are less than 28 days in age. 	
Key Points	 Silicone-based, latex-free, alcohol-free solution formulated to dissolve adhesives, (e.g., acrylic-based, rubber-based), and hydrocolloid-based residues from the skin. Product does not causing stinging as the product does not contain alcohol; may be used on broken/denuded skin. Product quickly evaporates. 	
Indications	 Use for all ages including neonates (see practice statement above): To assist with the removal of adhesive products, (e.g., tapes, ostomy flanges, hydrocolloids). To remove adhesive product residue from skin. 	
Precautions	 Extremely flammable; keep away for sources of ignition. Discontinue use if irritation noted. Do no use on around eyes, ears nose, mouth or vagina. 	
Contraindications	Do not use directly on wounds, burns or stomas.	
Formats & Sizes	 Wipe. Spray. 	

Directions	Rationale / Key Points	
Application to Remove a Product		
Do not open the package until ready to use. Ensure area is dry.	Opening the package immediately on use will ensure the wipe does not dry out.	
Gently lift a corner of the product to be removed and wipe or spray the adhesive remover between the skin and the product. Slowly peel back the product while continuing to wipe.		
Cleanse the area to remove any adhesive residue with warm dampened cloth/wipe or normal saline/sterile water-soaked gauze. Pat to dry the area.		
Ensure skin is dry before applying tape, ostomy flange, etc.		
Application to Remove Adhesive Residue		
Gently wipe or lightly spay the adhesive remover over area.		
Cleanse the area to remove any adhesive residue with warm		
dampened cloth/wipe or normal saline/sterile water-soaked gauze. Pat to dry the area.		
Ensure skin is dry before applying tape, ostomy flange, etc.		
Expected Outcomes		
Adhesive will be removed leaving the skin healthy and intact.	If product does not perform as expected, notify NSWOC/Wound Clinician and then consider submitting a <u>Supply Chain Product Concern Form</u> .	
For further information please contact NSWOC/Wound Clinician		