

Please refer to the individual Product Information Sheets for addition information.

Page 1/2

Skin Cleanser - Rinse	Skin Cleanser – No Rinse	Moisturize	Protectant Silicone 24%	Protectant Zinc 30%
Remedy Cleanse Shampoo & Body Wash 	Remedy Cleanse No Rinse 	Remedy Moisturize 	Remedy Prevent 	Secura Extra Protective Cream (EPC) 

For neonates in NICUs with a corrected gestational age greater than 31 weeks + 6 days or in maternity/pediatric units who are less than 28 days in age








<p>Indications: Used with a water bath for cleansing of skin and hair.</p> <p>Application:</p> <ul style="list-style-type: none"> Use a quarter size dab/dollop; gel will only produce a small amount of suds. Wash infant’s hair and body as per the neonatal unit’s practice/procedure for a water bath. Rinse the skin and hair. Pat dry. Do not apply moisturizer. If needed, apply a silicone protectant, (e.g., Remedy Prevent) to perineal area. <p>Note: Gel produces a small amount of suds as it is sulfate-free.</p> <p>Frequency of Use:</p> <ul style="list-style-type: none"> As needed. 	<p>Indications: For cleansing of skin and hair without water.</p> <p>Application:</p> <ul style="list-style-type: none"> Wash infant’s hair and body as per neonatal unit’s practice/procedure for sponge bath. Dispense 1-3 pumps onto hand or a warm damp cloth. Wash baby. Rinse if needed. Pat skin dry. Do not apply moisturizer. If needed, apply silicone protectant, (e.g., Remedy Prevent) to perineal area. <p>Perineal Cleansing:</p> <ul style="list-style-type: none"> Following each incontinence episode, cleanse area as per above. <p>Removal of Feces or Zinc:</p> <ul style="list-style-type: none"> Pump foam directly onto area. Allow to soak for 2-3 minutes. Gently wipe with a damp cloth to remove soiling or zinc. Repeat if needed. Do not rinse, pat skin dry. <p>Frequency of Use:</p> <ul style="list-style-type: none"> As needed. 	<p>Indications: As needed for treatment of very dry/cracked skin. DO NOT use for prevention of diaper dermatitis.</p> <p>Application:</p> <ul style="list-style-type: none"> Apply a <u>thin</u> layer to clean skin. Most effective when applied to slightly damp skin immediately after a bath. <p>Caution: DO NOT apply between toes. Remove any cream that does get between the toes.</p> <p>Frequency of Use:</p> <ul style="list-style-type: none"> Apply twice per day. 	<p>Indications:</p> <ul style="list-style-type: none"> Prevention of diaper dermatitis. Treatment of mild-moderate diaper dermatitis Prevention or treatment of moisture associated skin damage (MASD) for wounds, tubes/drains. <p>Application:</p> <ul style="list-style-type: none"> Apply a thin layer. If applied around wound, tube/drain, let dry 1-2 min before applying dressing. Do not use with silicone dressings. <p>Frequency of Use:</p> <ul style="list-style-type: none"> Prevention & treatment of Mild-Moderate Diaper Dermatitis: With each diaper change. Prevention & Treatment of Peri-wound/ tube/ drain MASD: Apply as ordered. 	<p>Indications:</p> <ul style="list-style-type: none"> Treatment of moderate-severe diaper dermatitis. Prevention with severe diarrhea, (e.g., Neonatal Abstinence Syndrome) <p>Application:</p> <ul style="list-style-type: none"> Apply cream onto palm of hand. <u>Apply in one direction.</u> The cream will “follow your hand”. Do not rub. Apply a 5mm thick layer to ensure <u>skin is not visible</u> through the cream. Repeat steps to cover affected area. Note: ostomy powder may be used over EPC to keep diaper zinc-free. <p>Cleansing an Incontinent Episode</p> <ul style="list-style-type: none"> Apply Remedy Cleanse No Rinse onto a damp cloth or directly onto soiled area. Wipe off the soiled EPC only. Re-apply more EPC cream as needed. Do not remove all EPC with every incontinence episode unless doing a skin assessment. <p>Frequency of Use:</p> <ul style="list-style-type: none"> After every incontinent episode. <p>Remove all EPC for skin assessment Q3days/prn:</p> <ul style="list-style-type: none"> Apply Remedy Cleanse No Rinse or the Shampoo & Body Wash (if doing a bath) directly onto the EPC. Let sit for 2-3 minutes then gently wipe with damp cloth. Repeat to remove all of the cream.
---	--	--	--	--

Neonatal Skin Care Protocol Additional Products

Wound Care Protocol

Please refer to the individual Product Information Sheets for addition information

Page 2/2

Barrier Film – Basic	Adhesive Remover	Barrier Film – Advanced	Skin Fold Transfer Sheets	Massage Oil	Hydrophilic Wound Dressing
Cavilon No-Sting Barrier Wipe 	Adapt Adhesive Remover 	Cavilon Advanced  or Marathon 	DriGo-HP  or Interdry 	Medium Chain Triglyceride Coconut Oil	Triad  To be used under the direction of an Wound/Ostomy Clinician or MPR.

For neonates in NICUs with a corrected gestational age greater than 31 weeks + 6 days or in maternity/pediatric units who are less than 28 days in age

<p>Indications: For intact or non-intact skin needing protection and/or additional securement, e.g.,</p> <ul style="list-style-type: none"> • Under dressings (IV, wound), ostomy flanges & tube/drain stabilization products. • Under medical devices. • On intact heels & elbows to minimize friction & shear damage. <p>Application</p> <ul style="list-style-type: none"> • Ensure area is clean. • Spray, or dab if using wipe, the film barrier over area and allow to dry. <p>Application for Crusting</p> <ul style="list-style-type: none"> • Apply a dusting of ostomy powder to the area. • Spray, or dab if using wipe, the film barrier over the powder and allow to dry. <p>If the area is weepy, repeat application of powder and film barrier to a maximum of 3 times; allow for the film barrier to dry between applications.</p>	<p>Indications:</p> <ul style="list-style-type: none"> • To assist in the removal of adhesive-based products, e.g., dressings, tapes, ostomy flanges, hydrocolloid • To remove residue from the skin due to adhesive products e.g. dressings, tapes, wafers, ostomy flanges, hydrocolloids. <p>Application</p> <ul style="list-style-type: none"> • Removal of adhesive product: <ul style="list-style-type: none"> ○ Gently lift a corner of the product; wipe adhesive remover between the skin and the product while gently peel back the product. ○ Cleanse area with water or normal saline & pat dry. • Removal of adhesive residue: <ul style="list-style-type: none"> ○ Apply remover, using wipe or light spray to the area. ○ Cleanse area with water or normal saline & pat dry. 	<p>Indications: To be used in consideration of the risks/benefits and under the direction of a NSWOC* or MRP when other skin protectants have not been effective in protecting the skin.</p>	<p>Indications: To be used in consideration of the risks/benefits and under the direction of a NSWOC or MRP for the treatment of intertrigo.</p>	<p>Indications: For weight gain and/or sensory stimulation and comfort.</p>	<p>Indications: • A zinc-based paste dressing used on open, moist/weeping areas & when left open to air, dries providing a cover-like dressing.</p> <p>Application:</p> <ul style="list-style-type: none"> • Apply a 3mm thick layer (thickness of a dime); wound should not be visible through the cream. • Position baby to allow cream to <u>dry for 15-20 mins</u>. If not able to be position, apply a gauze dressing overtop the paste to keep the paste in contact with the wound, not the baby's diaper. • Caution: Triad can be wiped off or washed off if in contact with fluid, (e.g., urine) if not allowed to dry or if not protected with gauze dressing. <p>Removal: Removal of paste is not needed with dressing changes but does need to be done prior to a Full Wound Assessment:</p> <ul style="list-style-type: none"> • Use sterile normal saline compress to wound area to soften the paste; leave in place 2-3 minutes then wipe off top layers. DO NOT rub. Repeat as needed. • See Product Information Sheet for other methods to remove the paste.
--	---	---	---	--	---

2025 March

All tubes and bottles are single patient use only; must be labelled with the baby's name and the date when opened.

*NSWOC = Nurse Specialized in Wound Ostomy Continence