Created by the British Columbia Provincial Professional Practice Stream – Wound Ostomy Continence









## **Product Information Sheet**

QuickChange Wrap			
Classification	Continence Containment Product / Male Penile Wrap		
Key Points	• Specially designed wrap made to encircle the penis; has a super -absorbent gel core that absorbs urine a way from the skin and an outer urine proof barrier to keep surrounding skin, linen and clothes dry.		
Indications	<ul> <li>Males who are bedfast, chair-fast or minimally mobile and require a small pad continence containment product that can manage up to 500ml of urine.</li> <li>May be used for those with a standard/longer-sized, small/retracted or hidden penis and those with scrotal edema.</li> <li>May be used with or without mesh pant.</li> </ul>		
Precautions	• For those with some mobility, use the wrap with a mesh pant.		
Contraindications	• Do not use with a brief, (e.g., pull-up, tabbed or belted).		
Formats & Sizes	• Wrap: • Onesize		

Directions	Rationale / Key Points
Selection	
For most applications, one wrap is most often used, but two wraps will be needed for a penis hidden by a large abdomen.	
Preparation	
Cleanse the peri-area. Apply skin protectant to a rea as per care plan.	
Application	
<ul> <li>For standard or long penis:</li> <li>Position the penis on the lower abdomen with penis facing straight towards the head.</li> <li>Slide wrap under the penis (Step1) with: <ul> <li>White side up</li> <li>Round edge towards the head</li> <li>The flaps (straight &amp; pointed) towards the legs.</li> </ul> </li> <li>Fold the pointed flap over top of the penis at a 45° angle (Step2); bring flap edge up to meet the rounded top edge.</li> <li>Fold the straight flap up over the wrap; slightly pull the flap upward to align its edge with the round edge (Step3).</li> <li>Using two fingers, pinch the hook fastener and inner pointed flap together. If needed, secure round edge with small piece of tape.</li> <li>Lay the wrap flat against the lower abdomen.</li> </ul>	
<ul> <li>For small or retracted penis:</li> <li>Slide wrap under the penis (Step1) with: <ul> <li>White side up</li> <li>Round edge towards the head</li> <li>The flaps (straight &amp; pointed) towards the legs.</li> </ul> </li> <li>Slide the pointed flap underneath the scrotum.</li> </ul>	

Adapted from Medline product information

Note: This is a controlled document; a printed copy may not reflect the current electronic version on the CLWK website (www.clwk.ca).

Created by the British Columbia Provincial Professional Practice Stream – Wound Ostomy Continence













## **Product Information Sheet**

Directions	Rationale / Key Points		
Lift the flap and scoop scrotum forward.			
<ul> <li>Pull the straight flap over and under the pointed flap (Step2).</li> <li>Using two fingers, pinch hook fastener and the inner pointed flap together. If needed, secure rounded edge with small piece of tape.</li> <li>If possible, lay the wrap on the lower abdomen (Step3.)</li> </ul>			
<ul> <li>For scrotal edema:</li> <li>Slide wrap under the penis (Step1) with: <ul> <li>White side up.</li> <li>Round edge towards the head.</li> <li>The flaps (straight &amp; pointed) towards the legs (Step2).</li> </ul> </li> <li>Slide the pointed flap underneath the scrotum.</li> <li>Pull the straight flap over and positon it under the pointed flap to create a bowl (Step3) which willcradle the edematous scrotum.</li> <li>Using two fingers, pinch the hook fastener and the inner pointed flap together.</li> </ul>			
<ul> <li>For hidden penis (need two wraps):</li> <li>If needed, lift abdominal skin-fold up to see peri-area.</li> <li>Slide the first wrap around scrotum (Step1) with: <ul> <li>White side up.</li> <li>Round edge towards the feet.</li> <li>The flaps (straight &amp; pointed) towards the head.</li> <li>Tuck the wrap between the thighs.</li> </ul> </li> <li>Place the second wrap (Step2): <ul> <li>White side up over the belly.</li> <li>Position the wrap to ensure the penis opening is visible.</li> <li>Fold edge of straight flap over to cover/hide fastener.</li> <li>Fold down the wrap lengthwise to cover the penis opening (Step3).</li> </ul> </li> <li>If needed, lay abdominals kin-fold down over the peri-area.</li> </ul>			
Removal			
Where one wrap is used; undo the fastener and remove wrap. Where two wraps have been used; remove the wraps.			
Frequency of Wrap Change			
Check wrap with every two hours to ensure wrap is in the correct position.	Client movement may move the wrap out of position		
Change wrap when wet.			
Expected Outcomes			
Urine is contained within the wrap protecting skin integrity.	If product does not perform as expected, notify NSWOC/		
Product performs as expected.	Wound Clinician and then consider submitting a <u>Supply</u> <u>Chain Product Concern</u> form.		
For further information please contact NSWOC/Wound Clinician			