




Product Information Sheet

Remedy Prevent	
Classification	Skin Care: Protectant - Silicone
Key Points	<ul style="list-style-type: none"> • 24% silicone blend with botanical nutrients providing a moisturizing protective barrier. • May be used on intact and non-intact skin. • Fragrance-free, hypoallergenic, paraben-free, sulfate-free. • Can be used in the presence of Chlorhexidine Gluconate (CHG). • Does not interfere with the integrity of latex, nitrile, and vinyl gloves.
Indications	<ul style="list-style-type: none"> • For all ages of patients/clients/residents for the: <ul style="list-style-type: none"> ○ Prevention and treatment of mild/moderate Incontinence Associated Dermatitis (IAD). ○ Prevention and treatment of Moisture Associated Skin Damage (MASD) around wounds, tubes/ drains. ○ Treatment of chafed, chapped or cracked skin.
Precautions	<ul style="list-style-type: none"> • When cream is used on the feet: <ul style="list-style-type: none"> ○ Do not apply cream between the toes; remove any cream that may inadvertently get between toes. ○ Ensure socks and shoes are worn when walking to prevent a fall.
Contraindications	<ul style="list-style-type: none"> • Do not use for clients with known sensitivity or allergy to product ingredients • Do not use on wounds e.g. Stage 2 Pressure Injury • Do not use on severe Incontinence-Associated Dermatitis (IAD) • Do not use with silicone-based dressings or tapes
Formats & Sizes	<ul style="list-style-type: none"> • Tube <ul style="list-style-type: none"> ○ 59mL ○ 256mL <div style="text-align: right;">  </div>

Directions	Rationale / Key Points
Selection	
Select the size of tube appropriate to the client situation, (e.g., short stay consider the 59mL sized tube). Label the tube with date and the client's name.	The tube is for single-patient use only and is to be used within three years of opening.
Application	
Apply a thin layer to clean skin and gently smooth in until cream is absorbed. Prevention of IAD/Treatment of mild/moderate IAD: Apply a thin layer to skin that may come in contact with urine, stool or moisture/exudate. Treatment of dry, chafed, chapped, cracked skin: <ul style="list-style-type: none"> • Apply a thin layer to dry, chafed, chapped or cracked skin. • For feet, do not apply cream between the toes; remove any cream that may inadvertently get between toes. Prevention/Treatment of Periwound, Peri-tube/drain MASD: Allow cream to dry for 1-2 minutes before apply a dressing. Note: cannot be used with silicone-based dressings or tapes.	Should not be applied so thick that there is a white layer and that the skin is not visible. A thin, transparent application provides good protection and allows for visual inspection of the skin. Product may make the soles of the feet slippery; ensure that socks/shoes are worn to prevent a fall. The silicone in the barrier cream may prevent the silicone cover dressings/tapes from adhering properly. This effect may also occur with other types of cover dressings/tapes



Product Information Sheet

Directions	Rationale / Key Points
Frequency of Application	
<p>After each incontinent episode, apply to clean and dry skin.</p> <p>Dry, chafed, chapped or cracked skin; reapply at least twice/day.</p>	
Expected Outcomes	
<p>Prevention of IAD.</p> <p>Resolution of mild /moderate IAD. If no resolution of IAD within 3 days, or condition worsen, then consider the use of a zinc-based cream and other continence strategies.</p> <p>Resolution of dry, chafed, chapped, cracked skin.</p> <p>Prevention/resolution of periwound, peri-tube/drain MASD.</p> <p>Product performs as expected.</p>	<p>If product does not perform as expected, notify NSWOC/Wound Clinician and then consider submitting a Supply Chain Product Concern Form.</p>
For further information please contact NSWOC/Wound Clinician	