1

🔊 fraser health



| Steri-Strips for Type 1 Skin Tears | | |
|----------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------|
| Classification | Adhesive Skin Closure Strip | |
| Key Points | • Strip is made with a porous, non-woven backing coated with a pressure-sensitive, hypoallergenic adhesive and reinforced with polyester filaments for added strength | |
| Indications | • Securing a viable flap with a Type 1 skin tear as defined by the International Skin Tear Advisory Panel | |
| Precautions | • Do not apply skin closures under tension; skin shearing, blistering or loss of adhesion may result if excessive tension is applied. | |
| Contraindications | When adhesion cannot be obtained, this indicated the presence of skin oils, moisture or hair. Infected wounds High tension wounds which cannot be easily approximated with gloved fingers. | |
| Formats & Sizes | Strips in various widths | |
| Ар | oplication Directions | Rationale |
| Ensure wound bed and flap have been cleansed and the flap has | | Must have optimal contact between the wound bed |
| been rolled back to cover the wound bed. | | and the flap. |
| Dry surrounding skin well. | | Skin must be dry for Steri-Strips to adhere. |
| Open Steri-Strip package and add the card to the aseptic field. | | |
| To Apply | | |
| Put on sterile gloves or use sterile forceps to handle the Steri- | | |
| Strips. | | |
| Start in the centre of the flap and without tension, apply the first | | Skin shearing, blistering or loss of adhesion my result if |
| Steri-Strip perpendicular to wound edge to hold flap in place. | | excessive tension is applied. |
| Secure one half of the Steri-Strip to the peri-wound skin, gently | | |
| approximate the wound edge together and laid the other half of | | |
| the Steri-Strip onto the opposite peri-wound skin area. Press | | |
| the Strip-Strip to the both peri-wound skin areas to secure it. | | |
| Secure the rest of the flap with the Steri-Strips, leaving a 1cm | | |
| space between each Steri-Strip for exudate to drain into cover | | |
| dressing. | | |
| | | |
| Steri-strips | | |
| Ē | | |
| Cover the re-approximated and secured flap with an | | The cover dressing is used to maintain moisture |
| appropriated cover dressing. | | balance of the wound bed to allow for optimal drainage management. |
| To Remove | | |
| DO NO REMOVE STERI-STRIPS; leave them to dry and fall off on | | Removal of Steri-strips may cause damage to the flap |
| their own. | | and surrounding skin. |
| Expected Outcome | | |
| | in flap securely in place. | |
| For further information, please contact your Wound Clinician. | | |
| | For further mormation, please con | nact your woulld Chiliciall. |

Providence